



Lie symmetry analysis and exact solutions of conformable time-space fractional Benjamin-Bona-Mahony equations

Amit^a, Rajeev Kumar^{a,*}, Anupma Bansal^b, Abdallah M. Talafha^c, Sandeep Malik^{d,**}

^aDepartment of Mathematics, M.M.E.C., Maharishi Markandeshwar Deemed to be University, Mullana (Ambala)-133001, Haryana, India

^bDepartment of Mathematics, D.A.V. College for Women, Ferozepur-152001, Punjab, India

^cDepartment of Mathematics and Natural Sciences, Prince Mohammad Bin Fahd University (PMU), AL Khobar, Saudi Arabia

^dDepartment of Mathematics, Akal University, Talwandi Sabo, Bathinda, 151302, Punjab, India

Abstract

This article investigates the invariance properties of nonlinear fractional partial differential equations involving conformable fractional derivatives in both time and space through the framework of Lie symmetry analysis. The study focuses on three significant forms of the Benjamin-Bona-Mahony (BBM) equations. The corresponding Lie symmetry generators and vector fields are systematically derived and employed to reduce the governing equations to ordinary differential equations. Exact analytical solutions are then constructed by utilising the (G'/G) -expansion method for the time-space fractional BBM equation. At the same time, suitable ansatz-based techniques are adopted to obtain explicit solutions for the modified and power-law fractional BBM equations. Furthermore, a qualitative dynamical analysis is carried out for the time-space fractional BBM equation and its modified form using bifurcation analysis and phase portraits, which reveal the behaviour of solution trajectories near the critical equilibrium points and provide insight into the nonlinear dynamics of the considered models. The results demonstrate that combining Lie symmetry analysis with analytical solution techniques provides an effective, systematic approach to studying nonlinear fractional BBM-type equations and deepens understanding of their mathematical structure and physical characteristics.

DOI: [10.46481/asr.2026.5.3.587](https://doi.org/10.46481/asr.2026.5.3.587)

Keywords: Benjamin-Bona-Mahony equation, Lie symmetry analysis, (G'/G) -expansion technique, Conformable fractional derivative, Bifurcation analysis

Article history:

Received: 17 June 2026

Received in revised form: 20 August 2026

Accepted for publication: 22 September 2026

Available online: 14 October 2026

© 2026 The Author(s). Published by the Nigerian Society of Physical Sciences under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

1. Introduction

Fractional calculus has attracted considerable attention over recent decades for addressing a variety of scientific problems, leading to significant theoretical advances. Several forms of fractional derivatives, including the Riemann–Liouville and Riesz derivatives, have been introduced to provide more accurate and realistic descriptions of physical processes. The Lie symmetry technique has also been extended to conformable differential equations for solving space-time fractional nonlinear partial differential equations

*Corresponding Author Tel. No.: +91-941-630-6632.

**Corresponding Author Tel. No.: +91-980-206-3796.

Email addresses: rajeevkumarbudhiraja@gmail.com (Rajeev Kumar), sndp796@gmail.com (Sandeep Malik)

(NLPDEs), while retaining derivative properties such as the chain rule. The conformable time and space fractional derivatives are defined, respectively, as follows [1–5]:

$$D^\beta[\Lambda(t)] = \lim_{\epsilon \rightarrow 0} \frac{\Lambda(t + \epsilon t^{1-\beta}) - \Lambda(t)}{\epsilon}, \quad (1)$$

$$D^\beta[\Lambda(x)] = \lim_{\epsilon \rightarrow 0} \frac{\Lambda(x + \epsilon x^{1-\beta}) - \Lambda(x)}{\epsilon}, \quad (2)$$

for each $t, x > 0$, where $\beta \in (0, 1]$. Suppose that the conformable fractional derivatives $D^\beta[\Lambda(t)]$ and $D^\beta[\Lambda(x)]$ are well defined for t and x , respectively, in $(0, b)$, with $b > 0$.

Researchers have developed numerous variants of the Benjamin–Bona–Mahony (BBM) equation because of its rich dynamics and broad range of applications. This well-known model describes the propagation of surface water waves in a uniform channel and has also been used for long-wavelength surface waves in liquids, acoustic waves in harmonic crystals, hydromagnetic waves in cold plasma, and acoustic-gravity waves in compressible fluids. This study focuses on three distinct time-space fractional BBM models.

The single-nonlinearity time-space fractional BBM equation is [6]:

$$T_t^\alpha(u) + uT_x^\alpha(u) - T_t^\alpha(T_x^\alpha(T_x^\alpha(u))) = 0. \quad (3)$$

The modified time-space fractional BBM equation is [6–9]:

$$T_t^\alpha(u) + u^2T_x^\alpha(u) - T_t^\alpha(T_x^\alpha(T_x^\alpha(u))) = 0. \quad (4)$$

The power-law time-space fractional BBM equation is [6, 10]:

$$T_t^\alpha(u) + u^nT_x^\alpha(u) - T_t^\alpha(T_x^\alpha(T_x^\alpha(u))) = 0. \quad (5)$$

Here, $u = u(x, t)$, and

$$T_t^\alpha(u) = t^{1-\alpha} \frac{\partial u}{\partial t}, \quad T_x^\alpha(u) = x^{1-\alpha} \frac{\partial u}{\partial x}, \quad (6)$$

$$T_t^\alpha(T_x^\alpha(T_x^\alpha(u))) = (1 - \alpha)t^{1-\alpha}x^{1-2\alpha} \frac{\partial^2 u}{\partial t \partial x} + t^{1-\alpha}x^{2-2\alpha} \frac{\partial^3 u}{\partial t \partial x^2}. \quad (7)$$

Also, $\alpha \in (0, 1]$.

Accordingly, effective analytical and numerical techniques have been developed to obtain solutions to these equations. Relevant methods include the exp-function scheme [11, 12], the Kudryashov method [13, 14], the sine–cosine method [15], the homotopy perturbation method [6], and the Riccati equation technique [16, 17]. Explicit solutions for these models have also been constructed using Lie symmetry methods and the (G'/G) -expansion method [18–21]. Numerical methods are likewise used to obtain approximate solutions when exact solutions are difficult or impossible to derive [22–24], providing insight into the dynamical behaviour of nonlinear models and complex physical phenomena.

The remainder of this paper is organised as follows. Section 2 determines the symmetries of the three equations using conformable fractional Lie symmetry analysis. Section 3 applies the (G'/G) -expansion and ansatz methods to obtain explicit exact solutions. Section 4 presents a bifurcation analysis of two reduced ordinary differential equations (ODEs), followed by the Conclusion.

2. Conformable fractional Lie symmetry analysis

First, the Lie group of point transformations is introduced [25–28]:

$$x^* = x + \epsilon \xi(x, t, u(x, t)) + O(\epsilon^2), \quad (8)$$

$$t^* = t + \epsilon \tau(x, t, u(x, t)) + O(\epsilon^2), \quad (9)$$

$$u^*(x^*, t^*) = u + \epsilon \eta(x, t, u(x, t)) + O(\epsilon^2). \quad (10)$$

where $\epsilon \ll 1$ is a free parameter. The Lie invariance criterion in prolonged form is

$$Pr^{(3)}V(E)|_{E=0} = 0, \quad (11)$$

which is appropriate for the highest-order derivatives appearing in the governing equation.

For completeness, the conformable fractional prolongation used to derive the determining equations is stated explicitly. The prolonged coefficients associated with the conformable derivatives are

$$\eta_t^\alpha = t^{1-\alpha} \eta_t + (1 - \alpha) \tau t^{-\alpha} u_t, \quad (12)$$

$$\eta_x^\alpha = x^{1-\alpha} \eta_x + (1-\alpha) \xi x^{-\alpha} u_x, \quad (13)$$

and

$$\begin{aligned} \eta_{xxt}^{3\alpha} = & (1-\alpha)(t^{1-\alpha} x^{1-2\alpha} \eta_{xt} + (1-2\alpha)t^{1-\alpha} x^{-2\alpha} \xi u_{xt} + (1-\alpha)t^{-\alpha} \tau x^{1-2\alpha} u_{xt}) \\ & + t^{1-\alpha} x^{2-2\alpha} \eta_{xxt} + (2-2\alpha)x^{1-2\alpha} t^{1-\alpha} \xi u_{xxt} + (1-\alpha)t^{-\alpha} x^{2-2\alpha} \tau u_{xxt}. \end{aligned} \quad (14)$$

Here, η_t , η_x , η_{xt} , and η_{xxt} denote the corresponding ordinary prolongation coefficients of the infinitesimal generator X , as defined in Section 3, Eqs. (11)–(12) of Refs. [27, 28]. Substitution of these ordinary prolongation coefficients into the conformable prolongation relations above gives the coefficients used in the present determining equations.

2.1. Single-nonlinearity time-space fractional BBM equation

Under the point transformations in Eqs. (8)–(10), u^* is a solution whenever u is a solution of Eq. (3). The invariance condition for Eq. (3) is

$$\eta_t^\alpha + u\eta_x^\alpha + u_x^\alpha \eta - \eta_{xxt}^{3\alpha} = 0. \quad (15)$$

Substituting the extended infinitesimals η , η_t^α , η_x^α , and $\eta_{xxt}^{3\alpha}$ and equating the coefficients of the resulting derivatives of u to zero yields the following overdetermined system of linear partial differential equations:

$$\xi_t = \xi_u = \tau_x = \tau_u = \eta_{uu} = \eta_{tu} = 0, \quad (16)$$

$$t^{1-\alpha} \eta_t + x^{1-\alpha} u \eta_x - (1-\alpha)t^{1-\alpha} x^{1-2\alpha} \eta_{xt} - t^{1-\alpha} x^{2-2\alpha} \eta_{xxt} = 0, \quad (17)$$

$$x^{1-\alpha} u \tau_t - (1-\alpha)x^{1-\alpha} t^{-1} u \tau + x^{1-\alpha} u \xi_x - (1-\alpha)x^{-\alpha} u \xi + x^{1-\alpha} \eta = 0, \quad (18)$$

$$2t^{1-\alpha} \xi_x - 2(1-\alpha)t^{1-\alpha} x^{-1} \xi - (1-\alpha)t^{1-\alpha} x^{1-2\alpha} \eta_{xu} - t^{1-\alpha} x^{2-2\alpha} \eta_{xxt} = 0, \quad (19)$$

$$t^{1-\alpha} x^{2-2\alpha} \xi_{xx} - (1-\alpha)t^{1-\alpha} x^{1-2\alpha} \xi_x + (1-\alpha)t^{1-\alpha} x^{-2\alpha} \xi - 2t^{1-\alpha} x^{2-2\alpha} \eta_{xu} = 0. \quad (20)$$

Solving this system gives the general solution

$$\eta = k_1 u, \quad \xi = k_2 x^{1-\alpha}, \quad \tau = -\frac{k_1}{\alpha} t + k_3 t^{1-\alpha}. \quad (21)$$

Here, k_1 , k_2 , and k_3 are arbitrary constants. The corresponding generators of the associated vector fields are

$$V_1 = -\frac{t}{\alpha} \frac{\partial}{\partial t} + u \frac{\partial}{\partial u}, \quad V_2 = x^{1-\alpha} \frac{\partial}{\partial x}, \quad V_3 = t^{1-\alpha} \frac{\partial}{\partial t}. \quad (22)$$

To derive the symmetry reduction of Eq. (3), the characteristic equation for the infinitesimal generator is

$$\frac{dx}{\xi} = \frac{dt}{\tau} = \frac{du}{\eta}, \quad (23)$$

where ξ , τ , and η are given by Eq. (21). Substituting the corresponding similarity solution into Eq. (3) yields the reduced ODE.

The vector field V_1 provides the following similarity variables:

$$\hat{\zeta} = x, \quad u = t^{-\alpha} H(\hat{\zeta}). \quad (24)$$

Substituting Eq. (24) into Eq. (3) gives the corresponding ODE

$$-\alpha H(\hat{\zeta}) + \hat{\zeta}^{1-\alpha} H(\hat{\zeta}) H'(\hat{\zeta}) + \alpha(1-\alpha) \hat{\zeta}^{1-2\alpha} H'(\hat{\zeta}) + \alpha \hat{\zeta}^{2-2\alpha} H''(\hat{\zeta}) = 0. \quad (25)$$

The vector field $V_2 + V_3$ provides the similarity variables

$$\hat{\zeta} = \frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha}, \quad u = H(\hat{\zeta}). \quad (26)$$

Substituting Eq. (26) into Eq. (3) gives

$$-H'(\hat{\zeta}) + H'(\hat{\zeta}) H(\hat{\zeta}) + H'''(\hat{\zeta}) = 0. \quad (27)$$

Integrating Eq. (27) with respect to $\hat{\zeta}$ introduces an arbitrary integration constant. Because this work focuses on localized travelling-wave solutions constructed using the (G'/G) -expansion method, the standard decay conditions $H, H', H'' \rightarrow 0$ as $|\hat{\zeta}| \rightarrow \infty$ are imposed. These conditions imply that the integration constant vanishes, $C = 0$, giving

$$2H''(\hat{\zeta}) + H(\hat{\zeta})^2 - 2H(\hat{\zeta}) = 0. \quad (28)$$

2.2. Modified time-space fractional BBM equation

Under the point transformations in Eqs. (8)–(10), u^* is a solution whenever u is a solution of Eq. (4). The invariance condition for Eq. (4) is

$$\eta_t^\alpha + u^2 \eta_x^\alpha + 2uu_x^\alpha \eta - \eta_{xxt}^{3\alpha} = 0. \quad (29)$$

Substituting the extended infinitesimals η_t^α , η_x^α , and $\eta_{xxt}^{3\alpha}$ and equating the coefficients of the resulting derivatives of u to zero yields the following overdetermined system of linear partial differential equations.

$$\xi_t = \xi_u = \tau_x = \tau_u = \eta_{uu} = \eta_{tu} = 0, \quad (30)$$

$$t^{1-\alpha} \eta_t + x^{1-\alpha} u^2 \eta_x - (1-\alpha) t^{1-\alpha} x^{1-2\alpha} \eta_{xt} - t^{1-\alpha} x^{2-2\alpha} \eta_{xxt} = 0, \quad (31)$$

$$2t^{1-\alpha} \xi_x - 2(1-\alpha) t^{1-\alpha} x^{-1} \xi - (1-\alpha) t^{1-\alpha} x^{1-2\alpha} \eta_{xu} - t^{1-\alpha} x^{2-2\alpha} \eta_{xxu} = 0, \quad (32)$$

$$x^{1-\alpha} u^2 \tau_t - (1-\alpha) x^{1-\alpha} t^{-1} u^2 \tau + x^{1-\alpha} u^2 \xi_x - (1-\alpha) x^{-\alpha} u^2 \xi + 2x^{1-\alpha} u \eta = 0, \quad (33)$$

$$t^{1-\alpha} x^{2-2\alpha} \xi_{xx} - (1-\alpha) t^{1-\alpha} x^{1-2\alpha} \xi_x + (1-\alpha) t^{1-\alpha} x^{-2\alpha} \xi - 2t^{1-\alpha} x^{2-2\alpha} \eta_{xu} = 0. \quad (34)$$

Solving this system gives

$$\eta = uk_1, \quad \xi = k_2 x^{1-\alpha}, \quad \tau = -\frac{2k_1}{\alpha} t + k_3 t^{1-\alpha}. \quad (35)$$

Here, k_1 , k_2 , and k_3 are arbitrary constants. The corresponding generators of the associated vector fields are

$$V_1 = -\frac{2t}{\alpha} \frac{\partial}{\partial t} + u \frac{\partial}{\partial u}, \quad V_2 = x^{1-\alpha} \frac{\partial}{\partial x}, \quad V_3 = t^{1-\alpha} \frac{\partial}{\partial t}. \quad (36)$$

To derive the symmetry reduction of Eq. (4), the characteristic equation for the infinitesimal generator is

$$\frac{dx}{\xi} = \frac{dt}{\tau} = \frac{du}{\eta}, \quad (37)$$

where ξ , τ , and η are given by Eq. (35). Substituting the solution associated with Eqs. (8)–(10) into Eq. (4) yields the reduced ODE.

The vector field V_1 gives the similarity variables

$$\hat{\xi} = x, \quad u = t^{-\frac{\alpha}{2}} H(\hat{\xi}). \quad (38)$$

Substituting Eq. (38) into Eq. (4) gives

$$-\frac{\alpha}{2} H(\hat{\xi}) + \hat{\xi}^{1-\alpha} H(\hat{\xi})^2 H'(\hat{\xi}) + \frac{\alpha(1-\alpha)}{2} \hat{\xi}^{1-2\alpha} H'(\hat{\xi}) + \frac{\alpha}{2} \hat{\xi}^{2-2\alpha} H''(\hat{\xi}) = 0. \quad (39)$$

The vector field $V_2 + V_3$ provides the similarity variables

$$\hat{\xi} = \frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha}, \quad u = H(\hat{\xi}). \quad (40)$$

Substituting Eq. (40) into Eq. (4) gives

$$-H'(\hat{\xi}) + H'(\hat{\xi}) H(\hat{\xi})^2 + H'''(\hat{\xi}) = 0. \quad (41)$$

Integrating Eq. (41) with respect to $\hat{\xi}$ introduces an arbitrary integration constant. Because this work focuses on localized homoclinic solutions constructed using the sech-expansion method, the standard decay conditions $H, H', H'' \rightarrow 0$ as $|\hat{\xi}| \rightarrow \infty$ are imposed. These conditions imply that the integration constant vanishes, $C = 0$, giving

$$3H''(\hat{\xi}) + H(\hat{\xi})^3 - 3H(\hat{\xi}) = 0. \quad (42)$$

2.3. Power-law time-space fractional BBM equation

Under the point transformations in Eqs. (8)–(10), u^* is a solution whenever u is a solution of Eq. (5). The invariance condition for Eq. (5) is

$$\eta_t^\alpha + u^n \eta_x^\alpha + nu^{n-1} u_x^\alpha \eta - \eta_{xxx}^{3\alpha} = 0. \quad (43)$$

Substituting the extended infinitesimals η_t^α , η_x^α , and $\eta_{xxx}^{3\alpha}$ and equating the coefficients of the resulting derivatives of u to zero yields the following overdetermined system of linear partial differential equations:

$$\xi_t = \xi_u = \tau_x = \tau_u = \eta_{uu} = \eta_{tu} = 0, \quad (44)$$

$$t^{1-\alpha} \eta_t + x^{1-\alpha} u^n \eta_x - (1-\alpha) t^{1-\alpha} x^{1-2\alpha} \eta_{xt} - t^{1-\alpha} x^{2-2\alpha} \eta_{xxx} = 0, \quad (45)$$

$$2t^{1-\alpha} \xi_x - 2(1-\alpha) t^{1-\alpha} x^{-1} \xi - (1-\alpha) t^{1-\alpha} x^{1-2\alpha} \eta_{xu} - t^{1-\alpha} x^{2-2\alpha} \eta_{xxu} = 0, \quad (46)$$

$$x^{1-\alpha} u^n \tau_t - (1-\alpha) x^{1-\alpha} t^{-1} u^n \tau + x^{1-\alpha} u^n \xi_x - (1-\alpha) x^{-\alpha} u^n \xi + nx^{1-\alpha} u^{n-1} \eta = 0, \quad (47)$$

$$t^{1-\alpha} x^{2-2\alpha} \xi_{xx} - (1-\alpha) t^{1-\alpha} x^{1-2\alpha} \xi_x + (1-\alpha) t^{1-\alpha} x^{-2\alpha} \xi - 2t^{1-\alpha} x^{2-2\alpha} \eta_{xu} = 0. \quad (48)$$

Solving this system gives

$$\eta = k_1 u, \quad \xi = k_2 x^{1-\alpha}, \quad \tau = -\frac{nk_1}{\alpha} t + k_3 t^{1-\alpha}. \quad (49)$$

Here, k_1 , k_2 , and k_3 are arbitrary constants. The corresponding generators of the associated vector fields are

$$V_1 = -\frac{nt}{\alpha} \frac{\partial}{\partial t} + u \frac{\partial}{\partial u}, \quad V_2 = x^{1-\alpha} \frac{\partial}{\partial x}, \quad V_3 = t^{1-\alpha} \frac{\partial}{\partial t}. \quad (50)$$

To derive the symmetry reduction of Eq. (5), the characteristic equation for the infinitesimal generator is

$$\frac{dx}{\xi} = \frac{dt}{\tau} = \frac{du}{\eta}, \quad (51)$$

where ξ , τ , and η are given by Eq. (49). Substituting the corresponding similarity solution into Eq. (5) yields the reduced ODE.

The vector field V_1 gives the following similarity variables:

$$\hat{\zeta} = x, \quad u = t^{-\frac{\alpha}{n}} H(\hat{\zeta}). \quad (52)$$

Substituting Eq. (52) into Eq. (5) gives

$$-\frac{\alpha}{n} H(\hat{\zeta}) + \hat{\zeta}^{1-\alpha} H(\hat{\zeta})^n H'(\hat{\zeta}) + \frac{\alpha(1-\alpha)}{n} \hat{\zeta}^{1-2\alpha} H'(\hat{\zeta}) + \frac{\alpha}{n} \hat{\zeta}^{2-2\alpha} H''(\hat{\zeta}) = 0. \quad (53)$$

The vector field $V_2 + V_3$ provides the similarity variables

$$\hat{\zeta} = \frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha}, \quad u = H(\hat{\zeta}). \quad (54)$$

Substituting Eq. (54) into Eq. (5) gives

$$-H'(\hat{\zeta}) + H'(\hat{\zeta}) H(\hat{\zeta})^n + H'''(\hat{\zeta}) = 0. \quad (55)$$

Integrating Eq. (55) with respect to $\hat{\zeta}$ introduces an arbitrary integration constant. Because this work focuses on localized homoclinic solutions constructed using the sech-expansion method, the standard decay conditions $H, H', H'' \rightarrow 0$ as $|\hat{\zeta}| \rightarrow \infty$ are imposed. These conditions imply that the integration constant vanishes, $C = 0$, giving

$$(n+1)H''(\hat{\zeta}) + H(\hat{\zeta})^{n+1} - (n+1)H(\hat{\zeta}) = 0. \quad (56)$$

3. Exact solutions of reduced ODEs

This section derives exact analytical solutions for the time-space fractional BBM models in Eqs. (3)–(5). The (G'/G) -expansion method [18–20] is used to obtain explicit solutions of Eq. (3). This method systematically constructs exact travelling-wave solutions by transforming the governing equation into a solvable ordinary differential equation.

For the modified time-space fractional BBM Eq. (4) and the power-law time-space fractional BBM Eq. (5), ansatz-based techniques are used to derive exact solutions. These methods are selected according to the nonlinear structure of the corresponding equations and permit the construction of analytical solutions. Together, the (G'/G) -expansion method and the ansatz approaches provide effective analytical techniques for investigating nonlinear fractional BBM-type equations.

3.1. Solution of Eq. (3)

To solve Eq. (28), assume that the ODE admits a solution expressed in powers of (G'/G) :

$$H(\hat{\zeta}) = e_m (G'/G)^m + e_{m-1} (G'/G)^{m-1} + \cdots + e_0, \quad (57)$$

where e_m are arbitrary constants, $m = 0, 1, 2, 3, \dots$, with $e_m \neq 0$. The function $G(\hat{\zeta})$ satisfies the second-order linear differential equation

$$G'' + \tilde{\omega}G' + \tilde{\delta}G = 0. \quad (58)$$

Here, $\tilde{\omega}$ and $\tilde{\delta}$ are arbitrary constants. For Eq. (28), take $H(\hat{\zeta}) \sim \hat{\zeta}^{-m}$, so that $H''(\hat{\zeta}) \sim \hat{\zeta}^{-m-2}$ and $H^2(\hat{\zeta}) \sim \hat{\zeta}^{-2m}$. Balancing the dominant terms H'' and H^2 gives $m + 2 = 2m$, and hence $m = 2$. Therefore, the ansatz in Eq. (57) becomes

$$H(\hat{\zeta}) = e_0 + e_1 (G'/G) + e_2 (G'/G)^2, \quad e_2 \neq 0. \quad (59)$$

From Eq. (59), we obtain

$$H(\hat{\zeta})^2 = e_0^2 + 2e_0e_1 (G'/G) + 2e_0e_2 (G'/G)^2 + e_1^2 (G'/G)^2 + 2e_1e_2 (G'/G)^3 + e_2^2 (G'/G)^4, \quad (60)$$

$$\begin{aligned} H''(\hat{\zeta}) &= (e_1\tilde{\omega}\tilde{\delta} + 2e_2\tilde{\delta}^2)(G'/G)^0 \\ &\quad + (e_1\tilde{\omega}^2 + 2e_1\tilde{\delta} + 6e_2\tilde{\delta}\tilde{\omega})(G'/G)^1 \\ &\quad + (3e_1\tilde{\omega} + 4e_2\tilde{\omega}^2 + 8e_2\tilde{\delta})(G'/G)^2 \\ &\quad + (2e_1 + 10e_2\tilde{\omega})(G'/G)^3 + 6e_2(G'/G)^4, \end{aligned} \quad (61)$$

Substituting Eq. (59) into Eq. (28) gives a polynomial in (G'/G) . Equating the coefficients of like powers of (G'/G) to zero yields

$$(G'/G)^4 : \frac{1}{2}e_2^2 + 6e_2, \quad (62)$$

$$(G'/G)^3 : e_1e_2 + 2e_1 + 10e_2\tilde{\omega}, \quad (63)$$

$$(G'/G)^2 : 3e_1\tilde{\omega} + e_0e_2 - e_2 + 8e_2\tilde{\delta} + 4e_2\tilde{\omega}^2 + \frac{1}{2}e_1^2, \quad (64)$$

$$(G'/G) : 6e_2\tilde{\delta}\tilde{\omega} + e_0e_1 + e_1\tilde{\omega}^2 + 2e_1\tilde{\delta} - e_1, \quad (65)$$

$$(G'/G)^0 : -e_0 + 2e_2\tilde{\delta}^2 + \frac{1}{2}e_0^2 + e_1\tilde{\omega}\tilde{\delta}, \quad (66)$$

Solving Eqs. (62)–(66) yields the following two parameter sets. For set 1,

$$\tilde{\delta} = \frac{1}{576}a^2 - \frac{1}{4}, \tilde{\omega} = -\frac{1}{12}a, e_0 = 3 - \frac{1}{48}a^2, e_1 = a, e_2 = -12. \quad (67)$$

For set 2,

$$\tilde{\delta} = \frac{1}{576}a^2 + \frac{1}{4}, \tilde{\omega} = -\frac{1}{12}a, e_0 = -1 - \frac{1}{48}a^2, e_1 = a, e_2 = -12. \quad (68)$$

Here, a is an arbitrary constant. Substituting set 1 into the ansatz in Eq. (59) gives

$$H(\hat{\zeta}) = 3 - \frac{1}{48}a^2 + a(G'/G) - 12(G'/G)^2, \quad (69)$$

For the exact solutions, let $q(\hat{\zeta}) = G'/G$, where G satisfies Eq. (58). For $\Delta = \tilde{\omega}^2 - 4\tilde{\delta} > 0$, the auxiliary equation has the solution

$$G(\hat{\zeta}) = e^{-\frac{\tilde{\omega}}{2}\hat{\zeta}} \left(J_1 \cosh\left(\frac{\sqrt{\Delta}}{2}\hat{\zeta}\right) + J_2 \sinh\left(\frac{\sqrt{\Delta}}{2}\hat{\zeta}\right) \right). \quad (70)$$

Hence,

$$\frac{G'}{G} = -\frac{\tilde{\omega}}{2} + \frac{\sqrt{\Delta}}{2} \frac{J_1 \sinh\left(\frac{\sqrt{\Delta}}{2}\hat{\zeta}\right) + J_2 \cosh\left(\frac{\sqrt{\Delta}}{2}\hat{\zeta}\right)}{J_1 \cosh\left(\frac{\sqrt{\Delta}}{2}\hat{\zeta}\right) + J_2 \sinh\left(\frac{\sqrt{\Delta}}{2}\hat{\zeta}\right)}. \quad (71)$$

The expanded solution is obtained as follows. If $\tilde{\omega}^2 - 4\tilde{\delta} > 0$, the exact solution of Eq. (3) is

$$u(x, t) = \left(\frac{a \sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(J_1 \sinh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \cosh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)}{2 \left(J_1 \cosh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \sinh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)} - \frac{\tilde{\omega}}{2} a \right)^2 - 12 \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(J_1 \sinh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \cosh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)}{2 \left(J_1 \cosh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \sinh \left(\frac{\sqrt{\tilde{\omega}^2 - 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)} - \frac{\tilde{\omega}}{2} \right)^2 + 3 - \frac{1}{48} a^2, \quad (72)$$

where J_1 and J_2 are arbitrary constants. Substituting set 2 into the ansatz in Eq. (59) gives

$$H(\hat{\xi}) = -1 - \frac{1}{48} a^2 + a(G'/G) - 12(G'/G)^2. \quad (73)$$

If $\tilde{\omega}^2 - 4\tilde{\delta} < 0$, the corresponding solution of Eq. (3) is

$$u(x, t) = \frac{a \sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(J_1 \sin \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \cos \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)}{2 \left(J_1 \cos \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \sin \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)} - \frac{\tilde{\omega}}{2} a \right)^2 + 12 \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(J_1 \sin \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \cos \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)}{2 \left(J_1 \cos \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) + J_2 \sin \left(\frac{\sqrt{-\tilde{\omega}^2 + 4\tilde{\delta}} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right)} \right) \right)} - \frac{\tilde{\omega}}{2} \right)^2 - 1 - \frac{1}{48} a^2, \quad (74)$$

where J_1 and J_2 are arbitrary constants.

Figure 1 shows the three-dimensional and corresponding two-dimensional representations of the kink-soliton solution $u(x, t)$. The solution exhibits a smooth, continuous transition between two distinct asymptotic states, and the two-dimensional profile illustrates its characteristic monotonic transition.

Figure 2 shows the three-dimensional and corresponding two-dimensional representations of the singular periodic solution $u(x, t)$. The solution exhibits a periodic wave structure containing singularities, while the two-dimensional profile shows the periodic oscillations and singular behaviour.

3.2. Solution of Eq. (4)

Assume that the ODE in Eq. (42) admits a solution of the form

$$H(\hat{\xi}) = A_0 + A_1 \operatorname{sech}(\kappa \hat{\xi}) + \cdots + A_p \operatorname{sech}(\kappa \hat{\xi})^p, \quad (75)$$

where A_i , $i = 0, 1, 2, \dots, p$, are arbitrary constants with $A_p \neq 0$. Balancing H'' and H^3 in Eq. (42) gives $p + 2 = 3p$, so $p = 1$. Therefore, the ansatz becomes

$$H(\hat{\xi}) = A_0 + A_1 \operatorname{sech}(\kappa \hat{\xi}). \quad (76)$$

Substituting Eq. (76) into Eq. (42) gives a polynomial in $\operatorname{sech}(\kappa \hat{\xi})$. Equating coefficients of like powers and solving the resulting algebraic system gives

$$A_0 = 0, \quad A_1 = \pm \sqrt{6}, \quad \kappa = \pm 1. \quad (77)$$

Substituting these values into Eq. (76) and then substituting the result into Eq. (40) gives the solution of Eq. (4):

$$u(x, t) = \pm \sqrt{6} \operatorname{sech} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha} \right). \quad (78)$$

Figure 3 shows the three-dimensional and corresponding two-dimensional representations of the bright-soliton solution $u(x, t)$. The surface and profile show a localised wave with a pronounced amplitude above the background level and decay away from the central peak, confirming the characteristic bright-soliton behaviour.




Figure 1: Kink-wave behaviour of the solution in Eq. (72) for $e_1 = -2$, $e_2 = -12$, $J_1 = 5$, $J_2 = 3$, and $\alpha = 0.5$.




Figure 2: Singular periodic-wave behaviour of the solution in Eq. (74) for $e_1 = 4$, $e_2 = -12$, $J_1 = 5$, $J_2 = 3$, and $\alpha = 1$.

3.3. Solution of Eq. (5)

Assume that the ODE in Eq. (56) admits a solution of the form

$$H(\hat{\zeta}) = A \operatorname{sech}(\kappa \hat{\zeta})^p, \quad (79)$$

where $A \neq 0$ is an arbitrary constant. In Eq. (56), the dominant terms are H'' and H^{n+1} . Thus, $p + 2 = (n + 1)p$, which gives $p = 2/n$.




Figure 3: Representation of the bright-soliton solution in Eq. (78) for $\alpha = 1$.

Therefore, the ansatz becomes

$$H(\hat{\zeta}) = A \operatorname{sech}\left(\kappa \hat{\zeta}\right)^{\frac{2}{n}}. \quad (80)$$

Substituting Eq. (80) into Eq. (56) gives a polynomial in sech . Equating coefficients of like powers and solving the resulting algebraic system gives

$$A = \left(\frac{(n+2)(n+1)}{2}\right)^{\frac{1}{n}}, \quad \kappa = \pm \frac{n}{2}. \quad (81)$$

Substituting these values into Eq. (80) and then substituting the result into Eq. (54) gives the solution of Eq. (5):

$$u(x, t) = \left(\frac{(n+2)(n+1)}{2}\right)^{\frac{1}{n}} \operatorname{sech}\left(\frac{n}{2} \left(\frac{x^\alpha}{\alpha} - \frac{t^\alpha}{\alpha}\right)\right)^{\frac{2}{n}}, \quad (82)$$

where n is a natural number.

Figure 4 shows the three-dimensional and corresponding two-dimensional representations of the bright-soliton solution $u(x, t)$. The solution forms a smooth, well-localised wave profile whose maximum amplitude is concentrated near the central region and decays towards the background in the far field. The two-dimensional profile further illustrates the localised nature and propagation characteristics of the bright soliton.

4. Bifurcation analysis of reduced ODEs

This section presents the bifurcation analysis [29–31] of the nonlinear autonomous systems associated with Eqs. (28) and (42). The critical points are identified, and their nature and stability are determined using the Jacobian matrix, eigenvalue analysis, and the corresponding first integral.

4.1. Bifurcation analysis corresponding to Eq. (28)

Eq. (28) can be expressed as the following nonlinear autonomous system:

$$H'(\hat{\zeta}) = G(\hat{\zeta}), \quad G'(\hat{\zeta}) = H(\hat{\zeta}) - \frac{H^2(\hat{\zeta})}{2}. \quad (83)$$

This nonlinear autonomous system has the critical points $(0, 0)$ and $(2, 0)$, which have different characteristics.




Figure 4: Representation of the bright-soliton solution in Eq. (82) for $\alpha = 1$ and $n = 3$.

At the critical point $(0, 0)$, the Jacobian matrix of the autonomous system is

$$\mathbf{J}(H, G) = \begin{pmatrix} 0 & 1 \\ 1 - H & 0 \end{pmatrix}. \quad (84)$$

Evaluating the Jacobian matrix at $(0, 0)$ gives

$$\mathbf{J}(0, 0) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (85)$$

The characteristic equation of the linearised system is

$$\lambda^2 - 1 = 0. \quad (86)$$

Thus, $\lambda_1 = -1$ and $\lambda_2 = 1$. The eigenvalues of the Jacobian matrix are real, distinct, and of opposite signs; therefore, the critical point $(0, 0)$ is an unstable saddle point, as shown in Figure 5.

At the critical point $(2, 0)$, introduce the shifted variables

$$h = H - 2, \quad g = G. \quad (87)$$

The original autonomous system is then transformed into

$$h'(\hat{\zeta}) = g(\hat{\zeta}), \quad g'(\hat{\zeta}) = -h(\hat{\zeta}) - \frac{h^2(\hat{\zeta})}{2}. \quad (88)$$

The Jacobian matrix of the transformed system at $(h, g) = (0, 0)$ is

$$\mathbf{J}(0, 0) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (89)$$

The characteristic equation of the linearised system is

$$\lambda^2 + 1 = 0. \quad (90)$$

Thus, $\lambda_{1,2} = \pm i$. The eigenvalues of the Jacobian matrix are purely imaginary. Because purely imaginary eigenvalues alone do not determine the stability of a nonlinear system, the linearisation is inconclusive. However, the system admits the first integral

$$G^2 - H^2 + \frac{H^3}{3} = C. \quad (91)$$

The level curves of this first integral in a neighbourhood of $(2, 0)$ are closed orbits. Therefore, the critical point $(2, 0)$ is a nonlinear centre; it is stable but not asymptotically stable, as shown in Figure 5.




Figure 5: Phase portrait of the nonlinear autonomous system in Eq. (83), showing the saddle point (0, 0) and the centre point (2, 0).

4.2. Bifurcation analysis corresponding to Eq. (42)

Eq. (42) can be expressed as the following nonlinear autonomous system:

$$H'(\hat{\zeta}) = G(\hat{\zeta}), \quad G'(\hat{\zeta}) = H(\hat{\zeta}) - \frac{H^3(\hat{\zeta})}{3}. \quad (92)$$

This nonlinear autonomous system has three critical points: (0, 0), ($\sqrt{3}$, 0), and ($-\sqrt{3}$, 0), with different characteristics.

At the critical point (0, 0), the Jacobian matrix of the autonomous system is

$$\mathbf{J}(H, G) = \begin{pmatrix} 0 & 1 \\ 1 - H^2 & 0 \end{pmatrix}. \quad (93)$$

Evaluating the Jacobian matrix at (0, 0) gives

$$\mathbf{J}(0, 0) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \quad (94)$$

The characteristic equation of the linearised system is

$$\lambda^2 - 1 = 0. \quad (95)$$

Thus, $\lambda_1 = -1$ and $\lambda_2 = 1$. The eigenvalues of the Jacobian matrix are real, distinct, and of opposite signs; therefore, the critical point (0, 0) is an unstable saddle point, as shown in Figure 6.

At the critical point ($\sqrt{3}$, 0), introduce the shifted variables

$$h = H - \sqrt{3}, \quad g = G. \quad (96)$$

The original autonomous system is then transformed into

$$h'(\hat{\zeta}) = g(\hat{\zeta}), \quad g'(\hat{\zeta}) = -2h(\hat{\zeta}) - \sqrt{3}h^2(\hat{\zeta}) - \frac{h^3(\hat{\zeta})}{3}. \quad (97)$$




Figure 6: Phase portrait of the nonlinear autonomous system in Eq. (92), showing the saddle point $(0,0)$ and the centre points $(\pm\sqrt{3},0)$.

The Jacobian matrix of the transformed system at $(h, g) = (0, 0)$ is

$$\mathbf{J}(0,0) = \begin{pmatrix} 0 & 1 \\ -2 & 0 \end{pmatrix}. \quad (98)$$

The characteristic equation of the linearised system is

$$\lambda^2 + 2 = 0. \quad (99)$$

Thus, $\lambda_{1,2} = \pm i\sqrt{2}$. The eigenvalues of the Jacobian matrix are purely imaginary. Because purely imaginary eigenvalues alone do not determine the stability of a nonlinear system, the linearisation is inconclusive. However, the system admits the first integral

$$G^2 - H^2 + \frac{H^4}{6} = C. \quad (100)$$

The level curves of this first integral in a neighbourhood of $(\sqrt{3},0)$ are closed orbits. Therefore, the critical point $(\sqrt{3},0)$ is a nonlinear centre; it is stable but not asymptotically stable, as shown in Figure 6.

At the critical point $(-\sqrt{3},0)$, introduce the shifted variables

$$h = H + \sqrt{3}, \quad g = G. \quad (101)$$

The original autonomous system is then transformed into

$$h'(\hat{\zeta}) = g(\hat{\zeta}), \quad g'(\hat{\zeta}) = -2h(\hat{\zeta}) + \sqrt{3}h^2(\hat{\zeta}) - \frac{h^3(\hat{\zeta})}{3}. \quad (102)$$

The Jacobian matrix of the transformed system at $(h, g) = (0, 0)$ is

$$\mathbf{J}(0,0) = \begin{pmatrix} 0 & 1 \\ -2 & 0 \end{pmatrix}. \quad (103)$$

The characteristic equation of the linearised system is

$$\lambda^2 + 2 = 0. \quad (104)$$

Thus, $\lambda_{1,2} = \pm i\sqrt{2}$. The eigenvalues of the Jacobian matrix are purely imaginary. Because purely imaginary eigenvalues alone do not determine the stability of a nonlinear system, the linearisation is inconclusive. However, the system admits the first integral

$$G^2 - H^2 + \frac{H^4}{6} = C. \quad (105)$$

The level curves of this first integral in a neighbourhood of $(-\sqrt{3}, 0)$ are closed orbits. Therefore, the critical point $(-\sqrt{3}, 0)$ is a nonlinear centre; it is stable but not asymptotically stable, as shown in Figure 6.

5. Conclusion

In this work, Lie symmetry analysis was applied to three forms of time-space conformable fractional BBM equations. The corresponding symmetry generators and vector fields were derived, demonstrating the effectiveness of the Lie group approach for conformable fractional partial differential equations. Using these symmetries, the governing equations were reduced to ODEs, from which exact trigonometric and hyperbolic solutions were obtained using the (G'/G) -expansion method and suitable ansatz techniques. Bifurcation analysis and phase portraits were also used to investigate the qualitative behaviour of two reduced systems, providing insight into the nature and stability of their solutions. The results show that combining Lie symmetry analysis with analytical solution methods provides an effective framework for studying nonlinear conformable fractional BBM-type equations.

Data availability

No new data were generated or analysed in this study. Data sharing is not applicable to this article.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this manuscript.

Funding

The authors received no specific funding from any public, commercial, or not-for-profit organisation for this research.

References

- [1] A. Akbulut & M. Kaplan, "Auxiliary equation method for time-fractional differential equations with conformable derivative", *Computers & Mathematics with Applications* **75** (2018) 876. <https://doi.org/10.1016/j.camwa.2017.10.016>.
- [2] T. Abdeljawad, "On conformable fractional calculus", *Journal of Computational and Applied Mathematics* **279** (2015) 57. <https://doi.org/10.1016/j.cam.2014.10.016>.
- [3] R. Khalil, M. Al-Horani, A. Yousef & M. Sababheh, "A new definition of fractional derivative", *Journal of Computational and Applied Mathematics* **264** (2014) 65. <https://doi.org/10.1016/j.cam.2014.01.002>.
- [4] D. R. Anderson & D. J. Ulness, "Newly defined conformable derivatives", *Advances in Dynamical Systems and Applications* **10** (2015) 109. Available online: <https://campus.mst.edu/adsa/contents/v10n2p2.pdf>.
- [5] R. Khalil, M. Al Horani & M. A. Hammad, "Geometric meaning of conformable derivative via fractional cords", *Journal of Mathematics and Computer Science* **19** (2019) 241. <https://doi.org/10.22436/jmcs.019.04.03>.
- [6] X. Sun & P. Yu, "Periodic traveling waves in a generalized BBM equation with weak backward diffusion and dissipation terms", *Discrete & Continuous Dynamical Systems-Series B* **24** (2019) 965. <https://doi.org/10.3934/dcdsb.2018341>.
- [7] M. A. Noor, K. I. Noor, A. Waheed & E. A. Al-Said, "Some new solitary solutions of the modified Benjamin-Bona-Mahony equation", *Computers & Mathematics with Applications* **62** (2011) 2126. <https://doi.org/10.1016/j.camwa.2011.06.060>.
- [8] K. Khan & M. A. Akbar, "Application of $\exp(-\phi(\xi))$ -expansion method to find the exact solutions of modified Benjamin-Bona-Mahony equation", *World Applied Sciences Journal* **24** (2013) 1373.
- [9] H. Naher & F. A. Abdullah, "The modified Benjamin-Bona-Mahony equation via the extended generalized Riccati equation mapping method", *Applied Mathematical Sciences* **6** (2012) 5495. Available online: <https://www.m-hikari.com/ams/ams-2012/ams-109-112-2012/naherAMS109-112-2012.pdf>.
- [10] A. G. Morgan, "Scattering for the quartic generalized Benjamin-Bona-Mahony equation", *Nonlinear Analysis* **261** (2025) 113909. <https://doi.org/10.1016/j.na.2025.113909>.
- [11] E. Hussain, A. H. Tedjani & M. A. S. Murad, "Exploring nonlinear dynamics of the (3+1)-dimensional Boussinesq-type equation: Wave patterns and sensitivity insight", *Axioms* **15** (2026) 198. <https://doi.org/10.3390/axioms15030198>.
- [12] S. Malik & S. Kumar, "Soliton solutions of dual-mode Kawahara equation via Lie symmetry analysis", in *Applied Analysis, Computation and Mathematical Modelling in Engineering*, S. S. Ray, H. Jafari, T. Raja Sekhar & S. Kayal (Eds.), Springer, Singapore, Singapore, 2022, pp. 189–202. https://doi.org/10.1007/978-981-19-1824-7_12.

- [13] M. A. S. Murad, "Optical solutions for perturbed conformable Fokas–Lenells equation via Kudryashov auxiliary equation method", *Modern Physics Letters B* **39** (2025) 2450418. <https://doi.org/10.1142/S0217984924504189>.
- [14] P. N. Ryabov, D. I. Sinelshchikov & M. B. Kochanov, "Application of the Kudryashov method for finding exact solutions of the high order nonlinear evolution equations", *Applied Mathematics and Computation* **218** (2011) 3965. <https://doi.org/10.1016/j.amc.2011.09.027>.
- [15] A. Bekir, "New solitons and periodic wave solutions for some nonlinear physical models by using the sine–cosine method", *Physica Scripta* **77** (2008) 045008. <https://doi.org/10.1088/0031-8949/77/04/045008>.
- [16] D. Zhuang, "A study of complex wave solutions and fractal structures for a (2+1) dimensional soliton system", *Far East Journal of Mathematical Sciences* **127** (2020) 71. <https://doi.org/10.17654/MS127010071>.
- [17] M. A. S. Murad, U. Younas, H. Emadifar, M. Iqbal, W. W. Mohammed & K. K. Ahmed, "Diverse soliton solutions to the conformable Ivancevic option pricing model via the modified generalized Riccati equation mapping method", *Mathematics and Computers in Simulation* **244** (2026) 213. <https://doi.org/10.1016/j.matcom.2025.12.022>.
- [18] A. M. Shahoot, K. A. E. Alurffi, M. O. M. Elmrif, A. M. Almsiri & A. M. H. Arwiniya, "The (G'/G) -expansion method for solving a nonlinear PDE describing the nonlinear low-pass electrical lines", *Journal of Taibah University for Science* **13** (2019) 63. <https://doi.org/10.1080/16583655.2018.1528663>.
- [19] E. M. E. Zayed, "The (G'/G) -expansion method combined with the Riccati equation for finding exact solutions of nonlinear PDEs", *Journal of Applied Mathematics & Informatics* **29** (2011) 351. Available online: <https://www.kci.go.kr/kciportal/ci/sereArticleSearch/ciSereArtiView.kci?sereArticleSearchBean.artiId=ART001520850>.
- [20] M. S. Hashemi & D. Baleanu, *Lie Symmetry Analysis of Fractional Differential Equations*, CRC Press, Boca Raton, FL, USA, 2020. <https://doi.org/10.1201/9781003008552>.
- [21] R. Kumar, S. Kumar, S. Kaur & S. Jain, "Time fractional generalized Korteweg–de Vries equation: Explicit series solutions and exact solutions", *Journal of Fractional Calculus and Nonlinear Systems* **2** (2021) 62. <https://doi.org/10.48185/jfcns.v2i2.315>.
- [22] E. V. Tikyaa, E. S. Nyiyongo, M. B. Ochang & S. D. Igba, "Simulation of optical fiber power amplification via higher-order soliton generation using the extended nonlinear Schrödinger equation", *Recent Advances in Natural Sciences* **3** (2025) 152. <https://doi.org/10.61298/rans.2025.3.1.152>.
- [23] A. A. Ishaq, F. M. Jimoh & K. Issa, "Numerical method for simulation of quadratic Riccati differential equations", *Recent Advances in Natural Sciences* **4** (2026) 159. <https://doi.org/10.61298/rans.2026.4.1.159>.
- [24] K. Issa, R. A. Bello & M. H. Sulaiman, "Approximate solution of time-fractional non-linear parabolic equations arising in mathematical physics", *African Scientific Reports* **3** (2024) 176. <https://doi.org/10.46481/asr.2024.3.2.176>.
- [25] S. Sahoo & S. S. Ray, "Lie symmetry analysis and exact solutions of (3+1) dimensional Yu–Toda–Sasa–Fukuyama equation in mathematical physics", *Computers & Mathematics with Applications* **73** (2017) 253. <https://doi.org/10.1016/j.camwa.2016.11.016>.
- [26] R. Kumar, R. Kumar & A. Bansal, "Lie symmetry and exact solutions of conformable time fractional Schamel–Korteweg–de Vries equation", *International Journal of Applied and Computational Mathematics* **10** (2024) 113. <https://doi.org/10.1007/s40819-024-01746-0>.
- [27] B. A. Tayyan & A. H. Sakka, "Lie symmetry analysis of some conformable fractional partial differential equations", *Arabian Journal of Mathematics* **9** (2020) 201. <https://doi.org/10.1007/s40065-018-0230-8>.
- [28] B. A. Tayyan & A. H. Sakka, "Symmetries and exact solutions of conformable fractional partial differential equations", *Palestine Journal of Mathematics* **9** (2020) 300. Available online: https://pjm.ppu.edu/sites/default/files/papers/PJM_October2019_300to311.pdf.
- [29] A. N. Ezaka, H. O. Adagba, S. N. Alope, L. O. Omenyi, T. E. Efor, O. Nwite & C. Agha, "Perturbation and bifurcation analysis of a primary-secondary dengue infection transmission model with explicit vector dynamics and optimal control", *African Scientific Reports* **5** (2026) 438. <https://doi.org/10.46481/asr.2026.5.2.438>.
- [30] Q. Chen, P. Kumar, Dipesh & H. M. Baskonus, "On the equilibrium point and Hopf-bifurcation analysis of GDP-national debt dynamics under the delayed external investment: A new DDE model", *Alexandria Engineering Journal* **91** (2024) 510. <https://doi.org/10.1016/j.aej.2024.02.022>.
- [31] F. M. Omar, M. A. S. Murad, S. S. Mahmood, S. Malik & T. Radwan, "Optical solitons, bifurcation, and chaos in the nonlinear conformable Schrödinger equation with group velocity dispersion coefficients and second-order spatiotemporal terms", *Scientific Reports* **15** (2025) 20073. <https://doi.org/10.1038/s41598-025-04387-5>.